

***Atrosynnema*, a new hyphomycete from dead branches in China**

Ji-WEN XIA, YING-RUI MA, JIAN-MEI GAO, XIU-GUO ZHANG & ZHUANG LI*

Department of Plant Pathology, Shandong Agricultural University, Taian, 271018, China

*CORRESPONDENCE TO: sdau613@163.com, liz552@126.com

ABSTRACT — *Atrosynnema digitosporum* gen. & sp. nov., collected on dead branches in Hainan Province, China, is described and illustrated. It has black synnematos conidiomata, integrated unbranched conidiophores, and monoblastic conidiogenous cells that produce brown, obclavate, dictyoseptate, rostrate conidia often with a mucilaginous cap around the apex.

KEY WORDS — conidial fungi, taxonomy

Introduction

Many new genera or species of saprobic darkly pigmented hyphomycetes continue to be discovered (e.g., Castañeda Ruíz et al. 2009, 2012, Cruz et al. 2012, Ma et al. 2014, Xia et al. 2014a,b). Hainan Province is considered an important reservoir of biodiversity, but its mycobiota is poorly known. We collected a fungus from dead branches that showed clear differences from all previously described asexual genera and is therefore described herein as a new genus (Ellis 1971, 1976; Subramanian 1971; Matsushima 1975, 1983, 1985, 1989, 1993, 1995; Carmichael et al. 1980; Castañeda-Ruiz 1986; Castañeda-Ruiz & Kendrick 1990a, b, 1991; Wu & Zhuang 2005; Seifert et al. 2011).

Samples were processed, examined and photographed following the methods described in Xia et al. (2014a). Adobe Photoshop 7.0 was used to process photographs into compound images and to modify backgrounds for esthetic reasons. Attempts were made to culture the species from single spores, but they did not germinate. Specimens are deposited in the Herbarium of the Department of Plant Pathology, Shandong Agricultural University, Taian,

China (HSAUP), and Mycological Herbarium, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China (HMAS).

***Atrosynnema* J.W. Xia, X.G. Zhang & Z. Li, gen. nov.**

MYCOBANK MB 816966

Differs from *Dictyospiropes* by synnematos conidiomata and monoblastic conidiogenous cells.

TYPE SPECIES: *Atrosynnema digitosporum* J.W. Xia et al.

ETYMOLOGY: *atro-* black, dark color; *-synnema* referring to the synnematos conidiomata.

ASEXUAL FUNGUS. COLONIES on the natural substrate effuse, brown. Mycelium immersed and superficial, of branched, septate, brown, smooth hyphae. CONIDIOMATA synnematos, unbranched, erect, with black stipes. CONIDIOPHORES integrated, erect, unbranched, septate, black. CONIDIOGENOUS CELLS monoblastic, integrated, black. Conidial secession schizolytic, with broad scars. CONIDIA solitary, dry, brown, smooth, dictyoseptate, obclavate, rostrate.

***Atrosynnema digitosporum* J.W. Xia, X.G. Zhang & Z. Li, sp. nov.**

FIG. 1

MYCOBANK MB 816967

Differs from *Dictyospiropes ziziphi* by synnematos conidiomata and monoblastic conidiogenous cells, producing smaller conidia with apices often surrounded by a globose mucilaginous cap.

TYPE: China, Hainan Province: Wuzhishan, dead branches of unidentified broadleaf tree, 22 Apr. 2015, J.W. Xia (Holotype, HSAUP H6651; isotype, HMAS 245596).

ETYMOLOGY: *digito-* finger-like rows or branches; *-sporum* referring to the conidia.

COLONIES on the natural substrate effuse, brown. Mycelium immersed and superficial, of branched, septate, brown, smooth, 2–4 µm wide hyphae. CONIDIOMATA synnematos, unbranched, erect, black, up to 350 µm long, up to 65 µm wide in the broadest part. CONIDIOPHORES integrated, erect, unbranched, septate, black, 6–10 µm wide. CONIDIOGENOUS CELLS monoblastic, cylindrical, integrated, black. Conidial secession schizolytic. CONIDIA solitary, dry, brown, smooth, dictyoseptate, obclavate, rostrate, 65–80 × 12–15.5 µm, apex often surrounded by a globose mucilaginous tunica.

COMMENTS – *Atrosynnema* resembles *Dictyospiropes* M.B. Ellis, but *Dictyospiropes* species have individual, polyblastic conidiogenous cells that produce obclavate, dictyosporous conidia without mucilaginous apices (Ellis 1976).

In conidial shape, *A. digitosporum* is most similar to *Dictyospiropes ziziphi* M.B. Ellis, but the conidia of *D. ziziphi* are larger, up to 140 µm long and 23 µm wide in the broadest part (Ellis 1976).




FIG. 1. *Atrosynnema digitosporum*. (ex holotype, HSAUP H6651): A. synnema, conidiophores, conidiogenous cells, and conidia; B. conidia.

Acknowledgments

The authors express gratitude to Dr. Bryce Kendrick and Dr. Rafael F. Castañeda-Ruiz for serving as pre-submission reviewers and for their valuable comments and suggestions. This project was supported by the National Natural Science Foundation of China (Nos. 31093440, 31230001) and the Ministry of Science and Technology of the People's Republic of China (Nos. 2006FY120100).

Literature cited

- Carmichael JW, Kendrick WB, Connors IL, Sigler YL. 1980. Genera of hyphomycetes. University of Alberta Press, Alberta. 386 p.
- Castañeda-Ruiz RF. 1986. Deuteromycotina de Cuba. Hyphomycetes IV. Instituto de Investigaciones Fundamentales en Agricultura Tropical "Alejandro de Humboldt", Habana, Cuba. 17 p.

- Castañeda-Ruiz RF, Kendrick WB. 1990a. Conidial fungi from Cuba: I. University of Waterloo Biology Series 32. 53 p.
- Castañeda-Ruiz RF, Kendrick WB. 1990b. Conidial fungi from Cuba: II. University of Waterloo Biology Series 33. 61 p.
- Castañeda-Ruiz RF, Kendrick WB. 1991. Ninety-nine conidial fungi from Cuba and three from Canada. University of Waterloo Biology Series 35. 132 p.
- Castañeda-Ruiz RF, Gusmão LFP, Cruz RAC da, Heredia Abarca G, Iturriaga T, Guarro J, Saikawa M, Stadler M, Minter DW. 2009. *Phaeocandelabrum*, a new genus of anamorphic fungi to accommodate *Sopagraha elegans* and two new species, *Ph. callisporum* and *Ph. joseiturriagae*. Mycotaxon 109: 221–232. <http://dx.doi.org/10.5248/109.221>
- Castañeda-Ruiz RF, Granados MM, Mardones M, Stadler M, Minter DW, Hernández-Restrepo M, Gené J, Guarro J. 2012. A microfungus from Costa Rica: *Ticosynnema* gen. nov.. Mycotaxon 122: 255–259. <http://dx.doi.org/10.5248/122.255>
- Cruz RAC da, Gusmão LFP, Castañeda-Ruiz RF, Stadler M, Minter DW. 2012. *Zelodactylaria*, an interesting new genus from semi-arid northeast Brazil. Mycotaxon 119: 241–248. <http://dx.doi.org/10.5248/119.241>
- Ellis MB. 1971. Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew. 608 p.
- Ellis MB. 1976. More dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew. 507 p.
- Ma LG, Xia JW, Ma YR, Castañeda-Ruiz RF, Zhang XG. 2014. Two new species of *Spadicoides* and *Gangliostilbe* from southern China. Mycological Progress 13: 547–552. <http://dx.doi.org/10.1007/s11557-013-0937-z>
- Matsushima T. 1975. Icones microfungorum a Matsushima lectorum. Kobe, Japan: Published by the author. 209 p.
- Matsushima T. 1983. Matsushima mycological memoirs 3. Published by the author, Kobe, Japan. 90 p.
- Matsushima T. 1985. Matsushima mycological memoirs 4. Published by the author, Kobe, Japan. 68 p.
- Matsushima T. 1989. Matsushima mycological memoirs 6. Published by the author, Kobe, Japan. 100 p.
- Matsushima T. 1993. Matsushima mycological memoirs 7. Published by the author, Kobe, Japan. 206 p.
- Matsushima T. 1995. Matsushima mycological memoirs 8. Published by the author, Kobe, Japan. 54 p.
- Seifert K, Morgan-Jones G, Gams W, Kendrick WB. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p.
- Subramanian CV. 1971. Hyphomycetes: an account of Indian species, except *Cercospora*. Indian Council of Agricultural Research, New Delhi. 930 p.
- Wu WP, Zhuang WY. 2005. *Sporidesmium*, *Endophragmiella* and related genera from China. Fungal Diversity Research Series 15. 351 p.
- Xia JW, Ma LG, Castañeda-Ruiz RF, Zhang XG. 2014a. *Minimelanolocus bicolorata* sp. nov., *Paradendryphiopsis elegans* sp. nov. and *Corynespora bannaense* sp. nov. from southern China. Mycoscience 55: 299–307. <http://dx.doi.org/10.1016/j.myc.2013.11.003>
- Xia JW, Ma LG, Castañeda-Ruiz RF, Zhang XG. 2014b. A new species of *Sporidesmiopsis* and three new records of other dematiaceous hyphomycetes from southern China. Nova Hedwigia 98: 103–111. <http://dx.doi.org/10.1127/0029-5035/2013/0145>